

Gravitomagnetism and spinor quantum mechanicsRonald J. Adler,^{1,2} Pisin Chen,^{3,4} and Elisa Varani⁵¹*Gravity Probe B, Hansen Laboratory for Experimental Physics, Stanford University, Stanford, California, 94305*²*Department of Physics, San Francisco State University, San Francisco California, 94132*³*Leung Center for Cosmology and Particle Astrophysics and Department of Physics and Graduate Institute of Astrophysics, National Taiwan University, Taipei, Taiwan 10617*⁴*Kavli Institute for Particle Astrophysics and Cosmology, SLAC National Accelerator Laboratory, Menlo Park California, 94025*⁵*Via Ponte Nuovo 24, 29014 Castell'Arquato (PC), Italy*

(Received 23 August 2011; published 26 January 2012)

We give a systematic treatment of a spin 1/2 particle in a combined electromagnetic field and a weak gravitational field that is produced by a slowly moving matter source. This paper continues previous work on a spin zero particle, but it is largely self-contained and may serve as an introduction to spinors in a Riemann space. The analysis is based on the Dirac equation expressed in generally covariant form and coupled minimally to the electromagnetic field. The restriction to a slowly moving matter source, such as the earth, allows us to describe the gravitational field by a gravitoelectric (Newtonian) potential and a gravitomagnetic (frame-dragging) vector potential, the existence of which has recently been experimentally verified. Our main interest is the coupling of the orbital and spin angular momenta of the particle to the gravitomagnetic field. Specifically, we calculate the gravitational g-factor to be $g_g = 1$; this is to be compared with the electromagnetic g-factor of $g_e = 2$ for a Dirac electron. Lastly, we discuss a number of possible experimental approaches to observing gravitomagnetic effects in atomic and macroscopic systems.

DOI: [10.1103/PhysRevD.85.025016](https://doi.org/10.1103/PhysRevD.85.025016)

PACS numbers: 03.65.-w, 03.65.Pm, 04.25.Nx, 04.80.Cc

I. INTRODUCTION

Classical systems in external gravitational fields have been studied for centuries, and recently the existence of the gravitomagnetic (or frame-dragging) field caused by the earth's rotation has been observed in two independent experiments. Observations of the LAGEOS satellites gave a measurement of the gravitomagnetic interaction via its effect on satellite orbits, accurate to about 10% [1]. The Gravity Probe B (GP-B) satellite verified the prediction of general relativity for the gravitomagnetic precession of a gyroscope in earth orbit (42 mas/yr) to about 20% [2–5]. Both experiments required impressive feats of data analysis and modeling of classical effects. Analysis of the LAGEOS data involved modeling the earth's Newtonian field to very high accuracy in order to extract the gravitomagnetic effect [6,7]. Analysis of the GP-B data required precise modeling of mechanical and electrical properties of the gyros [5,8]. It should be emphasized that the LAGEOS gravitomagnetic effects are due to Lorentz-like forces from the geodesic equation, whereas the GP-B gravitomagnetic effects are due to Larmor-like torques from the spin precession equation, so the two experiments are independent and rather nicely complementary.

While gravitomagnetic effects are generally quite small in the solar system, it is widely believed that they may play a large role in jets from active galactic nuclei so their experimental verification is of more than theoretical interest [9]. One can get a feel for the relative magnitude of gravitomagnetic effects using dimensional arguments to

see why satellite and laboratory experiments are so difficult, whereas for astrophysical sources such effects may be large. Gravitational effects for a system of mass M and size R generally involve a dimensionless factor of GM/c^2R (the metric distortion or the Newtonian potential divided by c^2), which for the earth is of order 10^{-9} . For gravitomagnetic effects there is an additional factor v_e/c for the spin velocity of the earth source, which is of order 10^{-6} . Finally, the gravitomagnetic force on a moving test body (similar to a Lorentz force) contains a factor v_{tb}/c , which is of order 10^{-5} for an orbiting satellite. Interestingly, the gravitomagnetic precession of a gyro is independent of the angular momentum of the rotor and thus also independent of the velocity due to its spin, as we will discuss in Sec. VIII. Thus, for experiments “near home” gravitomagnetic effects are generally much smaller than Newtonian effects. Contrariwise, for astrophysical systems containing neutron stars or black holes or active galactic nuclei all of the above numerical factors can be of order unity.

At the other end of the interest spectrum extensive theoretical work has been done on quantum fields in classical background spaces, the most well-known being related to Hawking radiation from black holes [2,10–12]. It is important to keep in mind that Hawking radiation has never been verified observationally.

Interesting experimental work has also been done on quantum systems in the earth's gravitational field, such as neutrons interacting with the earth's Newtonian field and atom interferometer experiments aimed at accurately testing the equivalence principle and other subtle general

relativistic effects [13–15]. There has been some discussion of attempts to see gravitomagnetic effects with these devices but such experiments would be quite difficult due to the small size of the effects and the similarity to classical effects of rotation; this is to be expected since gravitomagnetism manifests itself in a way that is similar to rotation, hence the appellation “frame-dragging.” Frame-dragging or “space-dragging” has been criticized as being technically inaccurate; the word gravitomagnetic is certainly more descriptive, albeit still awkward [16]. Laboratory detection of gravitomagnetic effects on a quantum system would clearly be of fundamental interest.

In this work we give a systematic treatment of a spin 1/2 particle in a combined electromagnetic field and weak gravitational field; this continues the work of reference [17]. We describe the particle with the generally covariant Dirac equation in a Riemann space, minimally coupled to the electromagnetic field in the standard gauge invariant way [18,19]. The weak gravitational field is naturally treated according to linearized general relativity theory, and we also assume a slowly moving matter source such as the earth [20–22]. Within this approximation the gravitational field is described by a gravitoelectric (or Newtonian) potential and a gravitomagnetic (or frame-dragging) vector potential, and the field equations are quite analogous to those of classical electromagnetism. We thus refer to it as the gravitoelectromagnetic (GEM) approximation. Our special emphasis throughout this paper is on the gravitomagnetic interaction.

The paper is organized as follows. After brief review comments on the GEM approximation (Sec. II) and the Dirac equation in flatspace (Sec. III), we give a detailed discussion of generally covariant spinor theory and the Dirac equation using the standard approach based on tetrads (Secs. IV and V) [19]. We then obtain the limit of the Dirac Lagrangian and the Dirac equation for a weak gravitational field, and discuss its interpretation in terms of an energy-momentum tensor (Sec. VI).

Our discussion of generally covariant spinors and the generally covariant Dirac equation is largely self-contained, and may serve as an introduction to the subject for uninitiated readers. In Sec. VI we also observe that the nongeometric or “flatspace gravity” approach of Feynman, Weinberg, and others does not appear to be completely equivalent to linearized general relativity theory in its coupling to spin [23]. We have not found this discussed elsewhere in the literature.

Using the weak gravitational field results, we then obtain the nonrelativistic limit of the theory (Sec. VII). We do this by integrating the interaction Lagrangian to obtain the interaction energy of the spinor particle with the electromagnetic and the GEM fields, and from that obtain the nonrelativistic interaction energies. This allows us to read off, in a simple and intuitive way, the interaction terms that one could use in a nonrelativistic Hamiltonian treatment. In

particular, we obtain (Sec. VIII) the usual g-factor of the electron $g_e = 2$ and the analogous result for the gravitomagnetic g-factor of a spinor, which is $g_g = 1$.

Section VIII also contains brief comments on the numerical value of some interesting and conceivably observable quantities, such as the precession of a spinning particle in the earth’s gravitomagnetic field and its relation to the precession of a macroscopic gyroscope; such precession appears to be universal for bodies with angular momentum. The phase shift in an atom interferometer is also mentioned as an experiment that could, in principle, show the existence of the gravitomagnetic field. Lastly, in Sec. VIII we mention that the acceleration of a body in a gravitational field depends on its angular momentum and estimate the small effect for atoms.

Finally, it is worth noting what we do *not* do in this paper. We study the effect of the gravitational field on a quantum mechanical spinor but not the effect of the spinor on the gravitational field or quantum gravity or quantum spacetime [24]. Similarly, we do not consider torsion, in which the affine connections have an antisymmetric part and are not equal to the Christoffel symbols. Some authors believe that inclusion of torsion is necessary for a full description of spin in general relativity [25]. However, Kleinert maintains that torsion is an alternative way to express the effects of curvature and the two are related by a novel sort of gauge transformation [26]. In any case, torsion has not proved necessary in our discussion.

II. THE GRAVITOELECTROMAGNETIC APPROXIMATION

In previous work we discussed linearized general relativity theory for slowly moving matter sources like the earth [17,21,22]. Here we summarize the results very briefly. The metric may be written as the Lorentz metric plus a small perturbation

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}. \quad (2.1)$$

We use coordinate freedom to impose the Lorentz gauge condition

$$\left(h_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} h \right)^{|\nu} = 0, \quad (2.2)$$

where the single slash denotes an ordinary derivative. Then, the field equations of general relativity tell us that the metric perturbation may be written as

$$h_{\mu\nu} = \begin{pmatrix} 2\phi & h^1 & h^2 & h^3 \\ h^1 & 2\phi & 0 & 0 \\ h^2 & 0 & 2\phi & 0 \\ h^3 & 0 & 0 & 2\phi \end{pmatrix}, \quad h_{00} = 2\phi, \quad h_{0k} = h^k, \quad (2.3)$$

where ϕ is the Newtonian or gravitoelectric potential and $h^k \leftrightarrow \vec{h}$ is the gravitomagnetic potential. For slowly moving sources, the field equations and the Lorentz condition become

$$\begin{aligned} \nabla^2 \phi &= 4\pi G\rho, & \nabla^2 h^j &= -16\pi G\rho v^j, \\ 4\dot{\phi} - \nabla \cdot \vec{h} &= 0, & \dot{\vec{h}} &= 0, \end{aligned} \quad (2.4)$$

where ρ is the source mass-energy density and v^j is its velocity.

The physical fields, which exert forces on particles, are the gravitoelectric (or Newtonian) field and the gravitomagnetic (or frame-dragging) field, which are defined by

$$\vec{g} = -\nabla\phi, \quad \vec{\Omega} = \nabla \times \vec{h}. \quad (2.5)$$

We call this equation system the GEM limit because of its close similarity to classical electromagnetism.

It is worth noting why the gravitational field in the GEM approximation is described by only the 4 metric components, ϕ and h^i , whereas gravity is generally described by 10 metric components. The energy-momentum tensor of

matter is $\rho u^\mu u^\nu$, where u^ν is the 4-velocity, so it is apparent that the spatial components are all of order v^2 . The field equations then imply that the spatial off-diagonal components of $h_{\mu\nu}$ are also of order v^2 and can be ignored for low source velocities, and also that the diagonal components are all equal to 2ϕ [17].

III. FLAT SPACE DIRAC EQUATION AND THE NONRELATIVISTIC LIMIT

We now briefly review the Dirac equation in flatspace and recast it into a Schrodinger equation form (SEF), which provides one convenient way to obtain the non-relativistic limit [18]. The SEF is exact and involves only the upper two components of the spinor wave function relevant for positive energy solutions in the nonrelativistic limit. This will serve as a basis of comparison for the alternative method we will use in Sec. VII for gravitational interactions. In this section, γ^μ denotes flatspace Dirac matrices [18,27].

The Dirac Lagrangian and the equations that follow from it are

$$L = a\bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi + b\bar{\psi}(-i\gamma^\mu \overleftarrow{\partial}_\mu - m)\psi - (a+b)eA_\mu \bar{\psi}\gamma^\mu \psi, \quad (3.1a)$$

$$(i\gamma^\mu \partial_\mu - m)\psi = eA_\mu \gamma^\mu \psi, \quad \bar{\psi}(-i\gamma^\mu \overleftarrow{\partial}_\mu - m) = eA_\mu \bar{\psi}\gamma^\mu. \quad (3.1b)$$

The spinor and its adjoint are considered independent in (3.1). The constants a and b are arbitrary, so long as $a+b \neq 0$. The γ^μ obey the flatspace Dirac algebra,

$$\{\gamma^\mu, \gamma^\alpha\} = 2\eta^{\mu\alpha}I. \quad (3.1c)$$

The adjoint spinor may be related to the spinor by a linear matrix relation, $\bar{\psi} = \psi^\dagger M$, where M is to be determined; consistency of the Eq. (3.1b) is assured if

$$M^{-1}\gamma^{\mu\dagger}M = \gamma^\mu, \quad M^{-1} = M = \gamma^0, \quad \bar{\psi} = \psi^\dagger \gamma^0. \quad (3.2)$$

Equation (3.2) is easy to verify for the choice of gamma matrices given below in (3.4).

The Hamiltonian form of the Dirac equation is gotten by multiplying (3.1) by γ^0 to obtain

$$\begin{aligned} i\partial_t \psi &= \beta m \psi + V + \vec{\alpha} \cdot \vec{\Pi} \psi, & \beta &\equiv \gamma^0, \\ \alpha &\equiv \gamma^0 \gamma^k, & \vec{p} &\equiv -i\nabla. \end{aligned} \quad (3.3)$$

Pauli's choice of gamma matrices is natural for the non-relativistic limit

$$\begin{aligned} \gamma^0 &= \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix}, & \gamma^i &= \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix}, \\ \vec{\alpha} &= \begin{pmatrix} 0 & \vec{\sigma} \\ \vec{\sigma} & 0 \end{pmatrix}. \end{aligned} \quad (3.4)$$

Next we break the 4-component wave function ψ into two 2-component Pauli spinors and factor out the time dependence due to the rest mass by substituting

$$\psi = e^{-imt} \begin{pmatrix} \Psi \\ \varphi \end{pmatrix}, \quad (3.5)$$

which leads to the coupled equations

$$\begin{aligned} i\partial_t \Psi &= V\Psi + (\vec{\sigma} \cdot \vec{\Pi})\varphi, \\ i\partial_t \varphi + 2m\varphi - V\varphi &= (\vec{\sigma} \cdot \vec{\Pi})\Psi. \end{aligned} \quad (3.6)$$

We are interested in Ψ so we solve for φ symbolically,

$$i\partial_t \Psi = V\Psi + (\vec{\sigma} \cdot \vec{\Pi})(2m - V + i\partial_t)^{-1}(\vec{\sigma} \cdot \vec{\Pi})\Psi, \quad (3.7a)$$

$$\varphi = (2m - V + i\partial_t)^{-1}(\vec{\sigma} \cdot \vec{\Pi})\Psi. \quad (3.7b)$$

The inverse operator $(2m - V + i\partial_t)^{-1}$ may be defined by its expansion in the time derivative, as discussed in Appendix A. The SEF (3.7a) is exact, although it is of infinite-order in the time derivative.

For the case of a free particle the operator factors on the right side of (3.7a) commute, and the equation becomes

$$i\partial_t \Psi = (i\partial_t + 2m)^{-1} \vec{p}^2 \Psi. \quad (3.8)$$

However the operators will not, in general, commute unless the field A_μ is constant.

In a low velocity system, the time variations of Ψ and V are associated with nonrelativistic energies, much less than the rest energy m , so we approximate the SEF (3.7a) by

$$i\partial_t\Psi = V\Psi + \frac{(\vec{\sigma} \cdot \vec{\Pi})^2}{2m}\Psi. \quad (3.9)$$

This is the spin 1/2 Schroedinger equation, often called the Pauli equation. It shows clearly how the spin and orbital angular momentum interact with the magnetic field. Pauli spin matrix algebra leads to an illuminating form for (3.9): to lowest-order in e ,

$$\begin{aligned} i\partial_t\Psi &= V\Psi + \frac{\vec{\Pi}^2}{2m}\Psi - \frac{e\vec{B} \cdot \vec{\sigma}}{2m}\Psi \\ &= V\Psi + \frac{\vec{p}^2}{2m}\Psi - \frac{e\vec{A} \cdot \vec{p}}{m}\Psi - \frac{e\vec{B} \cdot \vec{\sigma}}{2m}\Psi, \end{aligned} \quad (3.10)$$

where we have used the Lorentz gauge in which $\nabla \cdot \vec{A} = -\dot{A}^0$ and assumed A^0 has negligible time dependence.

The g-factor of a particle or system is defined in terms of its magnetic moment $\vec{\mu}$ and angular momentum \vec{J} by $\vec{\mu} = g_e(e/2m)\vec{J}$; thus, from (3.10), the fact that the energy is $-\vec{\mu} \cdot \vec{B}$ and the electron spin is $\vec{S} = \sigma/2$, it is evident that the electron g-factor is $g_e = 2$.

The relative coupling of the spin and orbital magnetic moments is made clearer if we consider a magnetic field that is approximately constant over the size of the system, in which case we can choose $\vec{A} = (\vec{B} \times \vec{r})/2$ and find from (3.10)

$$\begin{aligned} i\partial_t\Psi &= V\Psi + \frac{\vec{p}^2}{2m}\Psi - \frac{e\vec{B}}{2m}(2\vec{S} + \vec{L})\Psi, \\ \vec{S} &= \vec{\sigma}/2, \\ \vec{L} &= \vec{r} \times \vec{p}. \end{aligned} \quad (3.11)$$

That is $g_e = 2$ for the electron spin and $g_e = 1$ for the orbital angular momentum. This is, of course, approximate since QED gives corrections to the g-factor; to one-loop approximation the result is the famous anomalous moment $(g_e - 2)/2 = \alpha/2\pi$ [18,27].

The SEF (3.7a) may be expanded to higher-order to study e.g., hyperfine structure in the hydrogen atom [28]. An important problem is that the wave function Ψ is the upper-half of the Dirac wave function, so the quantity to be normalized is $|\Psi|^2 + |\varphi|^2$ rather than $|\psi_s|^2$ for a Schroedinger or Pauli wave function. To conserve probability, one must renormalize the wave function as discussed in detail in Ref. [28]. For this reason, that we will adopt an alternative and conceptually simpler approach to the non-relativistic limit in Sec. VII.

IV. GENERALLY COVARIANT SPINOR THEORY

The gravitational interaction of a spinor may be obtained most easily by making the Dirac Lagrangian (3.1a) and

Dirac Eq. (3.1b) generally covariant. We adopt the standard approach of using a tetrad of basis vectors in order to relate the generally covariant theory to the special relativistic theory in Lorentz coordinates [15,19]. This is a natural approach since Dirac spinors transform by the lowest dimensional representation S of the Lorentz group; that is $\psi' = S\psi$.

Two properties of the Dirac Lagrangian and Dirac equation must be modified to obtain a generally covariant theory: the Dirac algebra in (3.2) must be made covariant and the derivative of the spinor in (3.1) must be made into a covariant derivative.

The Dirac algebra (3.1c) is easily made covariant by replacing the Lorentz metric $\eta^{\mu\nu}$ by the Riemannian metric $g_{\mu\nu}$,

$$\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}I. \quad (4.1)$$

A set of γ^μ matrices that satisfy (4.1) is constructed by using a set of constant $\hat{\gamma}^b$ that satisfies the special relativistic relation (3.2) and a tetrad field e_b^μ normalized by the usual tetrad relations

$$e_b^\mu e_a^\nu g_{\mu\nu} = \eta_{ab}, \quad g^{\alpha\beta} = e_c^\alpha e_d^\beta \eta^{cd}. \quad (4.2)$$

Here the Greek indices label components of the tetrad vectors and Latin indices label the vectors. In terms of a convenient set of constant Dirac matrices $\hat{\gamma}^b$, such as those in (3.4), we define the γ^μ by

$$\gamma^\mu = e_b^\mu \hat{\gamma}^b. \quad (4.3)$$

It then follows from (3.1c) and (4.2) that the γ^μ satisfy

$$\{\gamma^\mu, \gamma^\nu\} = e_b^\mu e_a^\nu \{\hat{\gamma}^b, \hat{\gamma}^a\} = e_b^\mu e_a^\nu 2\eta^{ab}I = 2g^{\mu\nu}I. \quad (4.4)$$

The covariant derivative of a spinor is defined so as to transform as a vector under general coordinate transformations and as a spinor under Lorentz transformation of the tetrad basis. As with the covariant derivative of a vector, we define a rule for transplanting a spinor from x to a nearby point $x + dx$,

$$\psi^*(x + dx) = \psi(x) - \Gamma_\mu \psi(x) dx^\mu. \quad (4.5)$$

The matrices Γ_a are variously called spin connections, affine spin connections, or Fock-Ivanenko coefficients. The covariant derivative is then defined in terms of the difference between the value of the spinor and the value it would have if transplanted to the nearby point. That is

$$\begin{aligned} \psi(x)_{||\nu} dx^\nu &= [\psi(x) + \psi(x)_{|\nu} dx^\nu] - [\psi(x) \\ &\quad - \Gamma_\nu(x) \psi(x) dx^\nu] = [\psi(x)_{|\nu} \\ &\quad + \Gamma_\nu(x) \psi(x)] dx^\nu, \\ \psi_{||\nu} &= \psi_{|\nu} + \Gamma_\nu \psi = (\partial_\nu + \Gamma_\nu) \psi \equiv D_\nu \psi. \end{aligned} \quad (4.6)$$

Here the double slash denotes a covariant derivative. Since the spinor covariant derivative must transform as a vector

under coordinate transformations and as a spinor under Lorentz transformations of the tetrad basis, we have

$$\psi'_{||\mu} = \frac{\partial x^\nu}{\partial x'^\mu} S \psi_{||\nu}. \quad (4.7)$$

It follows from (4.6) and (4.7) that the spin connections must transform according to

$$\Gamma'_\nu = \frac{\partial x^\nu}{\partial x'^\mu} [S \Gamma_\nu S^{-1} - S_{|\nu} S^{-1}]. \quad (4.8)$$

The transformation (4.8) is formally similar to that of the Christoffel symbols, the affine connections used for vector covariant derivatives.

The covariant derivative of an adjoint spinor follows easily from that of a spinor in (4.6); we ask that the inner product $\bar{\psi}\chi$ of a spinor χ and an adjoint spinor $\bar{\psi}$ be a scalar and thus have a covariant derivative $(\bar{\psi}\chi)_{||\mu}$ equal to the ordinary derivative $(\bar{\psi}\chi)_{|\mu}$, and we also ask that the product rule hold for both the ordinary and the covariant derivatives. The result is

$$\bar{\psi}_{||\mu} = \bar{\psi}_{|\mu} - \bar{\psi}\Gamma_\mu. \quad (4.9)$$

The same idea leads to the covariant derivative of a gamma matrix; we ask that the expression $(\bar{\psi}\gamma^\mu\chi)_{||\alpha}$ be a second rank tensor and that it obey the product rule of differentiation, and find from (4.6) and (4.9)

$$\gamma^\mu_{||\omega} = \gamma^\mu_{|\omega} + \left\{ \begin{matrix} \mu \\ \omega\sigma \end{matrix} \right\} \gamma^\sigma + [\Gamma_\omega, \gamma^\mu]. \quad (4.10)$$

This expression plays an important role in obtaining the spin connections in the next section.

V. COVARIANT DIRAC LAGRANGIAN AND DIRAC EQUATION

In this section, we give a covariant Lagrangian and obtain the covariant Dirac equation. We also get a relation between the spinor and its adjoint and evaluate the spin connections.

The choice of a covariant Dirac Lagrangian L , and its associated density \mathcal{L} , is rather obvious from the flatspace Lagrangian in (3.1),

$$\begin{aligned} L &= a \bar{\psi}(i\gamma^\mu \psi_{||\mu} - m\psi) + b(-i\bar{\psi}_{||\mu} \gamma^\mu - \bar{\psi}m)\psi, \\ \mathcal{L} &= \sqrt{g}L. \end{aligned} \quad (5.1)$$

Coupling to the electromagnetic field will be included later. The γ^μ denotes the *covariant* Dirac matrices (4.3) throughout this section. The Dirac equations for the spinor and the adjoint spinor follow directly as the Euler-Lagrange equations of the Lagrangian density \mathcal{L} with ψ and $\bar{\psi}$ treated as independent variables,

$$(a + b)(i\gamma^\mu \psi_{||\mu} - m\psi) + ib\gamma^\mu_{||\mu} \psi = 0 \quad (5.2a)$$

$$(a + b)(\bar{\psi}_{||\mu} i\gamma^\mu + m\bar{\psi}) + ia\bar{\psi}\gamma^\mu_{||\mu} = 0. \quad (5.2b)$$

The spin connections, unspecified up to this point, may be chosen so that the divergence of γ^α vanishes, $\gamma^\mu_{||\mu} = 0$. The covariant Dirac equation is then the obvious generalization of the flatspace Eq. (3.1). The spin connections will be obtained below. Also, for symmetry and later convenience, we choose $a = b = 1/2$.

Next, as in flatspace in Sec. III, we ask that there be a relation between the adjoint and the spinor, $\bar{\psi} = \psi^\dagger M$, such that (5.2a) and (5.2b) are consistent. Manipulating (5.2a) we get for the adjoint,

$$\begin{aligned} -i\bar{\psi}_{|\mu} \tilde{\gamma}^\mu - i\bar{\psi} M_{|\mu}^{-1} M \tilde{\gamma}^\mu - i\bar{\psi} \tilde{\Gamma}_\mu \tilde{\gamma}^\mu - \bar{\psi}m &= 0, \\ \tilde{\gamma}^\mu &\equiv M^{-1} \gamma^{\mu\dagger} M, \quad \tilde{\Gamma}_\mu \equiv M^{-1} \Gamma_\mu^\dagger M. \end{aligned} \quad (5.3)$$

We then compare (5.3) with (5.2b), written as

$$-i\bar{\psi}_{|\mu} \gamma^\mu + i\bar{\psi} \Gamma_\mu \gamma^\mu - \bar{\psi}m = 0, \quad (5.4)$$

and see that M must satisfy the following two equations:

$$\gamma^\mu = \tilde{\gamma}^\mu = M^{-1} \gamma^{\mu\dagger} M, \quad (5.5a)$$

$$-\Gamma_\mu = \tilde{\Gamma}_\mu = M^{-1} \Gamma_\mu^\dagger M + M_{|\mu}^{-1} M. \quad (5.5b)$$

Equation (5.5a) may be written in terms of flatspace $\hat{\gamma}^b$ as

$$e_b^\mu \hat{\gamma}^b = e_b^\mu M^{-1} \hat{\gamma}^{b\dagger} M. \quad (5.6)$$

Thus, it is obvious that we should ask $\hat{\gamma}^b = M^{-1} \hat{\gamma}^{b\dagger} M$, as in flatspace (3.2), so we choose $M^{-1} = M = \hat{\gamma}^0$. Then, the derivative of M is zero, and it is easy to verify that the choice $M^{-1} = M = \hat{\gamma}^0$ also satisfies (5.5b).

Our remaining task is to obtain spin connections Γ_α . To do this, we make the natural demand that γ^μ have a null covariant derivative, so from (4.10)

$$\gamma^\mu_{||\alpha} = \gamma^\mu_{|\alpha} + \left\{ \begin{matrix} \mu \\ \alpha\beta \end{matrix} \right\} \gamma^\beta + [\Gamma_\alpha, \gamma^\mu] = 0. \quad (5.7)$$

This guarantees that the divergence vanishes, $\gamma^\mu_{||\mu} = 0$, as we have already mentioned. However, it is a stronger demand analogous to the demand that the metric have a null covariant derivative, which forces the affine connections to be Christoffel symbols. Note also that Γ_α is obviously arbitrary up to a multiple of the identity, which we will suppress henceforth.

To solve (5.7), we express γ^μ in terms of flatspace gammas $\hat{\gamma}^b$ as in (4.3) and rewrite (5.7) as

$$e_{b||\alpha}^\mu \hat{\gamma}^b + [\Gamma_\alpha, \hat{\gamma}^b] e_b^\mu = 0. \quad (5.8)$$

Multiplying this by the inverse tetrad matrix we get

$$[\Gamma_\alpha, \hat{\gamma}^c] = -e_\mu^c e_{b||\alpha}^\mu \hat{\gamma}^b. \quad (5.9)$$

We next note the well-known commutation relation on the sigma matrices, which are defined as $\hat{\sigma}^{ab} \equiv (i/2)[\hat{\gamma}^a, \hat{\gamma}^b]$,

$$[\hat{\sigma}^{ab}, \hat{\gamma}^c] = 2i(\hat{\gamma}^a \eta^{bc} - \hat{\gamma}^b \eta^{ac}). \quad (5.10)$$

From (5.10) it is evident that we should seek a solution that is proportional to $\hat{\sigma}^{ab}$ times a product of the tetrad and its derivatives. It is easy to verify that the choice

$$\Gamma_\alpha = \frac{i}{4} e_{b\mu} e_{a||\alpha}^\mu \hat{\sigma}^{ab}, \quad (5.11)$$

satisfies (5.9) and thus serves as the spin connection.

We thus have obtained a generally covariant theory in which the Lagrangian, the Dirac equations, the relation of the spinor to its adjoint, and the spin connections are generally covariant and consistent.

Finally, we include coupling to the electromagnetic field via minimal coupling, that is by substituting $iD_\mu \rightarrow iD_\mu - eA_\mu$; this gives the complete covariant Lagrangian

$$L = \frac{1}{2} \bar{\psi} (i\gamma^\mu \psi_{||\mu} - m\psi) + \frac{1}{2} (-i\bar{\psi}_{||\mu} \gamma^\mu - \bar{\psi}m)\psi - eA_\mu \bar{\psi} \gamma^\mu \psi. \quad (5.12)$$

We will study the weak gravitational field limit of this in the next section.

VI. LINEARIZED THEORY FOR WEAK GRAVITY

In this section, we use the results of Sec. V for covariant spinor theory to work out the weak field linearized theory. This is done by setting up an appropriate tetrad and using it to expand the Lagrangian (5.12) to lowest-order in the metric perturbation. The result is that there are three interaction terms in the Lagrangian, the first associated with the spin coefficients and the second with the alteration in the γ^μ caused by gravity. Remarkably the first vanishes in the linearized theory, while the second corresponds to an interaction via the energy-momentum tensor, as intuition should suggest. The third term is a cross term between the weak gravity and electromagnetic fields.

In a space with a nearly Lorentz metric (2.1), it is natural to choose a tetrad that lies nearly along the coordinate axes

$$e_a^\mu = \delta_a^\mu + w_a^\mu, \quad e_\nu^b = \delta_\nu^b - w_\nu^b, \quad (6.1)$$

where w_a^μ is a small quantity to be determined. From the fundamental tetrad relation (4.2) it follows that we should choose $w_{\mu\nu} = -(1/2)h_{\mu\nu}$ and thus have a tetrad and γ^μ matrices given by

$$e_a^\mu = \delta_a^\mu - (1/2)h_a^\mu, \quad (6.2)$$

$$\gamma^\mu = [\delta_a^\mu - (1/2)h_a^\mu] \hat{\gamma}^a = \hat{\gamma}^\mu - (1/2)h_a^\mu \hat{\gamma}^a.$$

Since Greek tensor indices and Latin tetrad indices are intimately mixed in the linearized theory, we will not distinguish between them in this section.

To evaluate the spin connections (5.11) with the tetrad (6.2), we need the Christoffel symbols and the covariant derivatives of the tetrad to first-order in $h_{\mu\nu}$,

$$\left\{ \begin{matrix} \nu \\ \mu\omega \end{matrix} \right\} = (1/2)(h_{\omega|\mu}{}^\nu + h_{\mu|\omega}{}^\nu - h_{\mu\omega}{}^{|\nu}), \quad (6.3)$$

$$e_{a||\mu}^\nu = (1/2)(h_{\mu|a}{}^\nu - h_{\mu a}{}^{|\nu}).$$

From (5.11), (6.2), and (6.3) we obtain the spin connections,

$$\Gamma_\mu = \frac{i}{4} e_{b\nu} e_{a||\mu}^\nu \hat{\sigma}^{ab} \cong \frac{i}{4} h_{\mu b|a} \hat{\sigma}^{ab}. \quad (6.4)$$

Thus, the Dirac Lagrangian (5.12) becomes

$$L = \frac{1}{2} \bar{\psi} (i\gamma^\mu \psi_{|\mu} - m\psi) + \frac{1}{2} (-i\bar{\psi}_{|\mu} \gamma^\mu - \bar{\psi}m)\psi - eA_\mu \bar{\psi} \gamma^\mu \psi + \frac{i}{2} \bar{\psi} \{ \hat{\gamma}, \Gamma_\mu \} \psi - \frac{i}{4} h_{\alpha}^\mu [\bar{\psi} \hat{\gamma}^\alpha \psi_{|\mu} - \bar{\psi}_{|\mu} \hat{\gamma}^\alpha \psi] + \frac{1}{2} h_{\alpha}^\mu A_\mu \bar{\psi} \hat{\gamma}^\alpha \psi, \quad (6.5)$$

with Γ_μ given in (6.4). The first line is the Dirac Lagrangian in flatspace (3.1a), and the other three terms are gravitational interactions that we now address.

The first interaction term in the second line of (6.5), due to the spin connections, contains the anticommutator $\{ \hat{\gamma}^\mu, \Gamma_\mu \}$. With the use of the symmetry of $h_{\mu\nu}$, the Dirac algebra (3.1c), and the operator identity $[AB, C] = A\{B, C\} - \{A, C\}B$ it is straightforward to verify the following two expressions:

$$h_{\mu b|a} \hat{\gamma}^\mu \hat{\sigma}^{ab} = i(h_{b|a}^a - h_{|b}^b) \hat{\gamma}^b, \quad (6.6)$$

$$h_{\mu b|a} \hat{\sigma}^{ab} \hat{\gamma}^\mu = i(h_{|b}^b - h_{b|a}^a) \hat{\gamma}^b,$$

and, thereby, see that

$$\{ \hat{\gamma}^\mu, \Gamma_\mu \} = \frac{i}{4} h_{\mu b|a} \{ \hat{\gamma}^\mu, \hat{\sigma}^{ab} \} = 0. \quad (6.7)$$

Thus, the interaction term containing the spin connections in (6.5) vanishes, which is a remarkable simplification. It should be stressed that this is only true to first-order, and the spin connections will generally be of interest in the nonlinear full theory.

There remains in the Lagrangian (6.5) only interactions due to the modification of the $\hat{\gamma}^\mu$ by gravity in (6.2); L may now be written as

$$L = \frac{1}{2} \bar{\psi} (i\hat{\gamma}^\mu \psi_{|\mu} - m\psi) + \frac{1}{2} (-i\bar{\psi}_{|\mu} \hat{\gamma}^\mu - \bar{\psi}m)\psi - eA_\mu \bar{\psi} \hat{\gamma}^\mu \psi - \frac{1}{4} h_{\alpha}^\mu [\bar{\psi} \hat{\gamma}^\alpha (i\psi_{|\mu} - eA_\mu \psi) - (i\bar{\psi}_{|\mu} + eA_\mu \bar{\psi}) \hat{\gamma}^\alpha \psi]. \quad (6.8)$$

The quantity in brackets in (6.8) is twice the appropriately symmetrized energy-momentum tensor $T^a{}_\mu$ for the Dirac field interacting with the electromagnetic field; that is, the gravitational interaction Lagrangian may be expressed as

$$L_{\text{IG}} = -\frac{1}{2}h_{\alpha}^{\mu}\left[\frac{1}{2}\bar{\psi}\hat{\gamma}^{\alpha}(i\psi_{|\mu} - eA_{\mu}\psi) - \frac{1}{2}(i\bar{\psi}_{|\mu} + eA_{\mu}\bar{\psi})\hat{\gamma}^{\alpha}\psi\right] = -\frac{1}{2}h_{\mu\alpha}T^{\mu\alpha}. \quad (6.9)$$

The energy-momentum tensor is discussed further in Appendix B.

The interaction (6.9) consists of the inner product of the field $h_{\mu\nu}$ with the conserved energy-momentum tensor $T^{\mu\nu}$; this coupling is in close analogy with the electromagnetic coupling between the field A_{μ} and the conserved current $j^{\mu} = e\bar{\psi}\gamma^{\mu}\psi$ in (6.8). Feynman has emphasized this analogy and developed a complete flatspace gravitational theory, with gravity treated as an ‘‘ordinary’’ two index (spin 2) field and formulated by analogy with electromagnetism, at least to lowest-order [23]. The geometric interpretation of gravity is, thereby, suppressed or ignored. Weinberg has similarly stressed that the geometric interpretation of gravity is not essential [15,23]. Schwinger also has used a similar and probably equivalent nongeometric methodology called source theory to obtain the standard results of general relativity theory, including the precession of a gyroscope due to the gravitomagnetic field [29]. However, there is a problem with relating the geometric and nongeometric viewpoints in that the Euler-Lagrange field equations are based on the Lagrangian *density* $\mathcal{L} = \sqrt{g}L \cong (1 + h/2)L$ and not the Lagrangian L , so there is an additional interaction term $(h/2)L$ in the geometric theory that is not present in the nongeometric theory; the equivalence of the Feynman approach to the linearized geometric approach is thus spoiled whenever the additional term does not vanish.

The difference between the Dirac equation per our geometric development and that which one would obtain from the nongeometric approach is easy to see. The Dirac equation that follows from (6.8) with $\mathcal{L} \cong (1 + h/2)L$ is

$$\gamma^{\mu}(i\psi_{|\mu} - eA_{\mu}) - m\psi = \frac{1}{2}h_{\mu\nu}\hat{\gamma}^{\mu}(i\psi^{|\nu} - eA^{\nu}\psi) + \frac{1}{4}(h^{\mu}{}_{\nu|\mu} - h_{|\nu})i\hat{\gamma}^{\nu}\psi. \quad (6.10)$$

The last term on the right containing $h_{|\nu}$ would not be present in the nongeometric approach. This will be discussed further in Sec. VII.

In summary of this section, the Lagrangian (6.8) contains the interaction of the Dirac field with the electromagnetic field to all orders and the interaction with the gravitational field only to lowest-order; (6.10) is the corresponding Dirac equation. We will discuss the interaction energies further in the following section in which we consider the nonrelativistic or low velocity limit of the theory.

VII. NONRELATIVISTIC LIMIT

We wish to use the results of the previous sections to obtain a nonrelativistic limit of the theory and calculate in a simple way some interesting properties of a spin 1/2 particle such as the electromagnetic g-factor and its gravitomagnetic analogue. The most familiar approach to this problem is to work with the upper two components of the Dirac wave function, as we did in Sec. III, and take the nonrelativistic limit [18,28]. However, the alternative approach we use in this section is conceptually simpler and avoids the problems of renormalization and Hermiticity that occur in the approach of Sec. III. The basic idea is to integrate the interaction Lagrangian over 3-space to get the interaction energy, then put the energy expression with Dirac 4-spinors into a form using Pauli 2-spinors, all in the low velocity limit [30].

In this section we will always work in nearly flatspace with Lorentz coordinates; the Dirac γ^{μ} will be those of flatspace and no hat will be used. Moreover, for simplicity, we will work in the Lorentz gauge for both the electromagnetic and GEM fields, and take both the Coulomb potential A^0 and the Newtonian potential ϕ to have negligible time dependence; that is $\dot{A}^0 = -\nabla \cdot \vec{A} = 0$ and $4\dot{\phi} = \nabla \cdot \vec{h} = 0$. This is appropriate for electromagnetic interactions in atoms and GEM interactions on the earth.

To illustrate the method we first consider only the electromagnetic interaction in flatspace; the results will be the same as those in Sec. III, in particular $g_e = 2$. The interaction Lagrangian and the interaction energy are, from (6.8),

$$L_{\text{IEM}} = -eA_{\mu}(\bar{\psi}\gamma^{\mu}\psi) = -A_{\mu}j^{\mu}, \quad (7.1a)$$

$$\Delta E_{\text{EM}} = -\int L_{\text{IEM}}d^3x. \quad (7.1b)$$

(I denotes interaction and EM electromagnetic.) For the Dirac ψ we use a convenient device, an expansion in terms of free positive energy Dirac wave functions on the mass shell. That is

$$\psi = \sum_{s=1,2} \int \frac{d^3p}{(2\pi)^3} f(p, s) [e^{ip_{\alpha}x^{\alpha}} u(p, s)], \quad (7.2)$$

$$E^2 = (p^0)^2 = \vec{p}^2 + m^2.$$

The positive energy wave functions do not form a complete set, but the approximation (7.2) should be quite good for distances much larger than the Compton wavelength, \hbar/m ; (7.2) is our fundamental assumption. A key idea in the calculation is to express the Dirac 4-spinor $u(p, s)$ in terms of a Pauli 2-spinor χ_s [30],

$$e^{-ip_{\alpha}x^{\alpha}} u(p, s) = e^{-ip_{\alpha}x^{\alpha}} \sqrt{\frac{E+M}{2m}} \left(\frac{I}{\frac{\vec{\sigma} \cdot \vec{p}}{E+M}} \right) \chi_s. \quad (7.3)$$

Correspondingly, we express the nonrelativistic Pauli wave function as

$$\Psi = \sum_{s=1,2} \int \frac{d^3 p}{(2\pi)^3} f(p, s) e^{ip_\alpha x^\alpha} \chi_s. \quad (7.4)$$

In terms of the above expressions (7.2) and (7.3), the interaction energy (7.1b) is

$$\begin{aligned} \Delta E_{\text{EM}} &= \sum_{s,s'=1,2} \int \frac{d^3 p}{(2\pi)^3} \frac{d^3 p'}{(2\pi)^3} f^*(p', s') f(p, s) \\ &\times \left[e \int d^3 x e^{i(p'_\alpha - p_\alpha) x^\alpha} \bar{u}(p', s') \gamma^\mu u(p, s) A_\mu \right]. \end{aligned} \quad (7.5)$$

The bracket in (7.5) corresponds to scattering of a free Dirac spinor by an external field, which is equivalent to scattering by an infinitely heavy source particle. It contains all the information about the spin interaction and corresponds to the diagram in Eq. (7.1) the particle leaves the wave function blob with 3-momentum \vec{p} , scatters from the external field via the QED vertex amplitude into momentum \vec{p}' , and then reenters the wave function blob. The electron remains on the mass shell, corresponding to zero energy transfer, which is consistent with a nonrelativistic wave function. We denote the 4-momentum transfer by $q_\mu = p'_\mu - p_\mu$, with $q_0 = 0$. The magnitude of the allowed 3-momentum transfer \vec{q} is limited by the width of the function $f(p, s)$ in momentum space.

It is now straightforward to calculate the bracket in (7.5). We split it into two parts, $\mu = 0$ for the electric interaction and $\mu = j$ for the magnetic interaction. For the electric part, we have

$$\begin{aligned} &e \int d^3 x e^{iq_\alpha x^\alpha} A_0 \bar{u}(p', s') \gamma^0 u(p, s) \\ &= e \int d^3 x e^{iq_\alpha x^\alpha} A_0 \left(\frac{E+m}{2m} \right) \chi_{s'}^\dagger \left[I, \frac{\vec{\sigma} \cdot \vec{p}'}{E+M} \right] \left[\frac{I}{E+m}, \frac{\vec{\sigma} \cdot \vec{p}}{E+m} \right] \chi_s \\ &= e \int d^3 x e^{iq_\alpha x^\alpha} A_0 \chi_{s'}^\dagger \left[\frac{E}{m} + \frac{\vec{q} \cdot \vec{p}}{2m(E+m)} + \frac{i\vec{q} \times \vec{p} \cdot \vec{\sigma}}{2m(E+m)} \right] \chi_s. \end{aligned} \quad (7.6)$$

The first term in the bracket in (7.6) is the obvious charge coupling to the Coulomb field. The second and third terms may be simplified. First, because there is no energy




FIG. 1. The electron in the wave function scatters from the field and back into the wave function.

transferred $\vec{p}^2 = \vec{p}'^2$, from which it follows that $\vec{p} \cdot \vec{q} = -\vec{q}^2/2$. Secondly, the vector \vec{q} multiplying the exponential may be replaced by $i\nabla$ operating on the exponential, after which integration by parts allows us to replace it with $-i\nabla$ operating on the function A_0 ; that is, we may replace $\vec{q}A_0 \rightarrow -i\nabla A_0$. Thus, the second term vanishes since $\nabla^2 A_0 = 0$ in a charge-free region for the Lorentz gauge. What remains is to order $1/m^2$,

$$\begin{aligned} &e \int d^3 x e^{iq_\alpha x^\alpha} A_0 \bar{u}(p', s') \gamma^0 u(p, s) \\ &= \int d^3 x e^{iq_\alpha x^\alpha} \left[e(\chi_{s'}^\dagger \chi_s) A_0 + \frac{e}{4m^2} \nabla A_0 \times \vec{p} \cdot (\chi_{s'}^\dagger \vec{\sigma} \chi_s) \right]. \end{aligned} \quad (7.7)$$

The second term in (7.7) is clearly a fine structure correction, which we mentioned in Sec. IV and which will not concern us further [28].

The $\mu = j$ magnetic part of the interaction in (7.5) is handled in exactly the same way as the electric part. We have

$$\begin{aligned} &e \int d^3 x e^{iq_\alpha x^\alpha} A_j \bar{u}(p', s') \gamma^j u(p, s) \\ &= e \int d^3 x e^{iq_\alpha x^\alpha} A_j \left(\frac{E+m}{2m} \right) \chi_{s'}^\dagger \left[I, \frac{\vec{\sigma} \cdot \vec{p}'}{E+m} \right] \\ &\quad \times \left[\begin{array}{cc} 0 & \sigma^j \\ \sigma^j & 0 \end{array} \right] \left[\frac{I}{E+m}, \frac{\vec{\sigma} \cdot \vec{p}}{E+m} \right] \chi_s \\ &= \int d^3 x e^{iq_\alpha x^\alpha} A_j \left(\frac{e}{2m} \right) \chi_{s'}^\dagger [\sigma^j \vec{\sigma} \cdot \vec{p} + \vec{\sigma} \cdot \vec{p}' \sigma^j] \chi_s \\ &= - \int d^3 x e^{iq_\alpha x^\alpha} \left(\frac{e}{2m} \right) \chi_{s'}^\dagger [2\vec{p} \cdot \vec{A} + \vec{q} \cdot \vec{A} + i\vec{q} \times \vec{A} \cdot \vec{\sigma}] \chi_s. \end{aligned} \quad (7.8)$$

We then replace $\vec{q} \rightarrow -i\nabla$ and see that the second term in the bracket vanishes in a gauge with $\nabla \cdot \vec{A} = 0$, so we are left with

$$\begin{aligned} &e \int d^3 x e^{iq_\alpha x^\alpha} A_j \bar{u}(p', s') \gamma^j u(p, s) \\ &= - \int d^3 x e^{iq_\alpha x^\alpha} \left(\frac{e}{2m} \right) \chi_{s'}^\dagger [2\vec{p} \cdot \vec{A} + \nabla \times \vec{A} \cdot \vec{\sigma}] \chi_s \\ &= - \int d^3 x e^{iq_\alpha x^\alpha} \left[\frac{e}{m} \vec{p} \cdot \vec{A} (\chi_{s'}^\dagger \chi_s) + \frac{e}{2m} \vec{B} \cdot (\chi_{s'}^\dagger \vec{\sigma} \chi_s) \right]. \end{aligned} \quad (7.9)$$

Finally, we combine (7.7) and (7.9) and substitute into (7.5) to obtain to order $1/m$,

$$\begin{aligned}\Delta E_{\text{EM}} &= \sum_{s,s'=1,2} \int \frac{d^3 p}{(2\pi)^3} \frac{d^3 p'}{(2\pi)^3} f^*(p', s') f(p, s) \\ &\quad \times \int d^3 x e^{-i(\vec{p}' - \vec{p}) \cdot \vec{x}} \chi_{s'}^\dagger \left[e A_0 - \frac{e}{m} \vec{p} \cdot \vec{A} - \frac{e}{2m} \vec{B} \cdot \vec{\sigma} \right] \chi_s \\ &= \int d^3 x \Psi^\dagger \left[e A_0 - \frac{e}{m} \vec{p} \cdot \vec{A} - \frac{e}{2m} \vec{B} \cdot \vec{\sigma} \right] \Psi. \quad (7.10)\end{aligned}$$

This is the same result that we obtained in (3.10) of Sec. III, so we have thus verified that our present approach reproduces the usual result for the electron g-factor, $g_e = 2$.

We now work out the nonrelativistic limit of the gravitational interaction in (6.8), following the same procedure as for the electromagnetic interaction; we will not include the product of the electromagnetic and gravitational fields, that is the cross term in (6.8). The algebra is a bit lengthier but equally straightforward. As with the Lagrangian and energy for the electromagnetic case in (7.1), we have for the gravitational case

$$L_{\text{IG}} = -\frac{1}{2} h_{\mu\nu} T^{\mu\nu}, \quad \Delta E_G = -\int L_{\text{IG}} d^3 x, \quad (7.11)$$

where $T^{\mu\nu}$ is given in (6.9), I denotes interaction, and G denotes gravity. It is convenient to write $T^{\mu\nu}$ in close analogy with the electromagnetic current as

$$T^{\mu\alpha} = \bar{\psi} \gamma^\alpha \left(\frac{1}{2} i \overleftrightarrow{\partial}^\mu \right) \psi. \quad (7.12)$$

Note the relation between the electromagnetic and the gravitational interactions,

$$A_\mu \leftrightarrow h_{\mu\nu}/2, \quad \gamma^\mu \leftrightarrow \gamma^\mu \left(\frac{i}{2} \overleftrightarrow{\partial}^\nu \right). \quad (7.13)$$

Then ΔE_G is, in analogy with (7.5),

$$\begin{aligned}\Delta E_G &= \sum_{s,s'=1,2} \int \frac{d^3 p}{(2\pi)^3} \frac{d^3 p'}{(2\pi)^3} f^*(p', s') f(p, s) (h_{\mu\nu}/2) \\ &\quad \times \int d^3 x e^{i(p'_\alpha - p_\alpha)x^\alpha} \bar{u}(p', s') \gamma^\mu \left(p^\nu + \frac{q^\nu}{2} \right) u(p, s). \quad (7.14)\end{aligned}$$

As with the electromagnetism calculation we split the gravitational interaction into two parts, the gravitoelectric for $h_{00} = h_{ii} = 2\phi$ and the gravitomagnetic for $h_{0j} = h_{j0} = h^j$. The gravitoelectric part of the bracket in (7.14) involves the same spin products as encountered with the electromagnetic calculation in (7.7) and (7.9), and after some algebra we obtain to order $1/m^2$,

$$\begin{aligned}&\left[\int d^3 x e^{i(p'_\alpha - p_\alpha)x^\alpha} \bar{u}(p', s') \left\{ \gamma^0 E + \left(p^j + \frac{q^j}{2} \right) \gamma^j \right\} u(p, s) \phi \right] \\ &= \int d^3 x e^{i(p'_\alpha - p_\alpha)x^\alpha} \chi_{s'}^\dagger \left[\left(\frac{E^2}{m} \phi + \frac{E}{4m^2} \nabla \phi \times \vec{p} \cdot \vec{\sigma} \right) \right. \\ &\quad \left. + \left(\frac{\vec{p}^2}{m} \phi + \frac{1}{2m} \nabla \phi \times \vec{p} \cdot \vec{\sigma} \right) \right] \chi_s \\ &= \int d^3 x e^{i(p'_\alpha - p_\alpha)x^\alpha} m \chi_{s'}^\dagger \left[\left(1 + \frac{2\vec{p}^2}{m^2} \right) \phi \right. \\ &\quad \left. + \frac{3}{4m^2} \nabla \phi \times \vec{p} \cdot \vec{\sigma} \right] \chi_s. \quad (7.15)\end{aligned}$$

A word is in order about the physical interpretation of the gravitoelectric result (7.15). The term $m\phi$ is of course the expected Newtonian energy; the factor $(1 + 2\vec{p}^2/m^2)$ occurs also in the analysis of a spin zero system in Ref. [17], and is approximately the Lorentz transformation factor between the potential in the lab frame and the moving frame of the particle; thus $(1 + 2\vec{p}^2/m^2)\phi$ is the Newtonian potential seen by the moving particle. The last term in the bracket has the same form and is the gravitational analog of the fine structure term in the electromagnetic energy (7.7), except of course for the different coefficient. We will not be concerned further with the higher-order terms in (7.15) and will, henceforth, keep only the lowest-order term ϕ in the bracket.

We turn finally to the gravitomagnetic part of the interaction (7.14), which is our main interest in this work. The gravitomagnetic part of the bracket, proportional to h^j , is

$$\begin{aligned}&\left[\int d^3 x e^{i(p'_\alpha - p_\alpha)x^\alpha} (h^j/2) \bar{u}(p', s') \left\{ \gamma^0 (p^j + q^j/2) \right. \right. \\ &\quad \left. \left. + E \gamma^j \right\} u(p, s) \right] \\ &= \left[\int d^3 x e^{i(p'_\alpha - p_\alpha)x^\alpha} u^\dagger(p', s') \left\{ (p^j + q^j/2) (h^j/2) \right. \right. \\ &\quad \left. \left. + E (h^j/2) \alpha^j \right\} u(p, s) \right]. \quad (7.16)\end{aligned}$$

Note that the term $\vec{q} \cdot \vec{h}$ will vanish by gauge choice, just as the $\vec{q} \cdot \vec{A}$ term vanished for the electromagnetic case. Then, using the same previous manipulations on the spin products, we reduce this to

$$\begin{aligned}&\left[\int d^3 x e^{i(p'_\alpha - p_\alpha)x^\alpha} \bar{u}(p', s') \left\{ \gamma^0 \left(p^j + \frac{q^j}{2} \right) + E \gamma^j \right\} u(p, s) \frac{h^j}{2} \right] \\ &= \left[\int d^3 x e^{i(p'_\alpha - p_\alpha)x^\alpha} \chi_{s'}^\dagger \left\{ \vec{p} \cdot \vec{h} + \frac{1}{4} \nabla \times \vec{h} \cdot \vec{\sigma} \right\} \chi_s \right], \quad (7.17)\end{aligned}$$

where we have neglected terms of higher-order, that is $1/m^2$. Finally, we combine (7.15) and (7.17) to obtain the total energy

$$\begin{aligned}
\Delta E_G &= \sum_{s,s'=1,2} \int \frac{d^3 p}{(2\pi)^3} \frac{d^3 p'}{(2\pi)^3} f^*(p', s') f(p, s) \\
&\quad \times \left[\int d^3 x e^{i(p'_\alpha - p_\alpha)x^\alpha} \chi_{s'}^\dagger \left(m\phi + \vec{p} \cdot \vec{h} \right. \right. \\
&\quad \left. \left. + \frac{1}{4} \nabla \times \vec{h} \cdot \vec{\sigma} \right) \chi \right] \\
&= \int d^3 x \Psi^\dagger \left(m\phi + \vec{p} \cdot \vec{h} + \frac{1}{4} \vec{\Omega} \cdot \vec{\sigma} \right) \Psi. \quad (7.18)
\end{aligned}$$

(Recall that the gravitomagnetic field is $\vec{\Omega} = \nabla \times \vec{h}$.) This is the main result of this section and is consistent with the result of Ref. [17] for a scalar particle.

Finally, we note that since we have expanded the wave function in terms of a free Dirac particle on the mass shell (7.2) the free Dirac Lagrangian is zero and the extra geometric interaction term $(\hbar/2)L$ discussed in Sec. VI vanishes.

VIII. GRAVITOMAGNETIC PHYSICAL EFFECTS

The result (7.18) is to be compared with the analogous electromagnetic result (7.10). We see, of course, that the Newtonian potential is the analog of the Coulomb potential eA^0 and the gravitomagnetic potential is the analog of the vector potential according to

$$eA^0 \leftrightarrow m\phi, \quad (-e/m)\vec{A} \leftrightarrow \vec{h}. \quad (8.1)$$

We also see that the coupling of the spin to the gravitomagnetic field $\vec{\Omega}$ is only half the analogous electromagnetic coupling. To make this most obvious we consider a gravitomagnetic field $\vec{\Omega}$ that is approximately constant over the system so that we may choose $\vec{h} = (\vec{\Omega} \times \vec{r})/2$. Then,

$$\begin{aligned}
\Delta E_G &= \int d^3 x \Psi^\dagger \left(m\phi + \frac{1}{2} \vec{\Omega} \times \vec{r} \cdot \vec{p} + \frac{1}{4} \vec{\Omega} \cdot \vec{\sigma} \right) \Psi \\
&= \int d^3 x \Psi^\dagger \left(m\phi + \frac{1}{2} \vec{\Omega} \cdot \vec{r} \times \vec{p} + \frac{1}{2} \vec{\Omega} \cdot \frac{\vec{\sigma}}{2} \right) \Psi \\
&= \int d^3 x \Psi^\dagger \left[m\phi + \frac{1}{2} \vec{\Omega} \cdot (\vec{L} + \vec{S}) \right] \Psi. \quad (8.2)
\end{aligned}$$

Orbital and spin angular momenta couple in the same way to the gravitomagnetic field, so the g-factor for gravitomagnetism is $g_g = 1$ for both.

Hehl and Ni have studied a Dirac equation in an accelerated and rotating frame and find a similar coupling of spin plus orbital angular momentum to the rotation rate ω , with the equivalence $\omega \leftrightarrow \Omega/2$ [31].

From the correspondence between the magnetic and gravitomagnetic couplings it is clear that since a magnetic moment due to orbital angular momentum, $(e/2m)\vec{L}$, precesses at the Larmor frequency $(eB/2m)$ in a magnetic field B , the gravitomagnetic moment due to both orbital and spin angular momenta will precess in a

gravitomagnetic field Ω with frequency $\Omega/2$. Thus, precession of a quantum system should be the same as that observed in the classical gyroscopes of the GP-B satellite experiment [5]. It thus seems very likely that the precession rate is universal for any angular momentum system, whether the angular momentum is classical or quantum mechanical, orbital, or spin.

For the surface of the earth the magnitude of the gravitomagnetic field is quite small, as estimated in Ref. [17]. The field and the associated quantum energy are of order

$$\Omega \approx 10^{-13} \text{ rad/s}, \quad E_\Omega = \hbar\Omega \approx 10^{-28} \text{ eV}. \quad (8.3)$$

Experimental detection of such small quantum gravitomagnetic effects in an earth-based lab would obviously be difficult. Such an experiment might be performed with an atomic interferometer. The atomic beam could be split into two components with angular momenta differing by $\Delta J \approx \hbar$. Then, according to (8.2) the two components would have energies differing by about $\Delta E \approx \Omega \Delta J \approx \Omega \hbar$ and thus suffer phase shifts differing by about $\Delta \varphi \approx \Delta E t / \hbar \approx \Omega t$, where t is the time of flight. For a typical $t = 1 \text{ s}$ this implies a phase shift of order 10^{-13} rad, which is orders of magnitude less than presently detectable [32].

In addition to the small size of gravitomagnetic effects one might see in the laboratory there is a serious further inherent difficulty in such experiments; a rotation of the apparatus would in general have similar effects, as noted above, and swamp the gravitomagnetic effects. Such rotations would have to be controlled and compensated to very high accuracy as mentioned in the introduction and in Refs. [17,31].

A different aspect of gravitomagnetism is the effect of angular momentum on the free fall motion of a body. It has long been recognized that, according to general relativity, bodies with internal structure or angular momentum do not exactly follow geodesics, and thus are not appropriate test bodies; detailed corrections to the motion have been calculated [20,33]. The effect on motion is easily seen in the present context from the coupling in (8.2) between the angular momentum $J = L + S$ and Ω ; for both quantum and classical systems, $\Delta E = J_k \Omega_k / 2$. If Ω is inhomogeneous, this implies a force on the system in exactly the same way that a magnetic moment feels a force in an inhomogeneous magnetic field [34]. Thus, for example, two atoms with opposite spin directions in the earth's field will undergo different accelerations. The difference is easy to estimate; the force is the gradient of the energy,

$$F_j = (1/2) \Omega_{k|j} J_k. \quad (8.4)$$

Thus, we may estimate roughly, using (8.3) and $J \approx \hbar$, that the acceleration difference for an atom of mass m_a is

$$\Delta g \cong \hbar \Omega / R_e m_a \cong 10^{-28} \text{ m/s}^2, \quad (8.5)$$

where R_e is the radius of the earth. As with the other effects we have discussed this is smaller (by about nine-orders of

magnitude) than has been even optimistically considered for testing the equivalence principle (EP) in earth orbit, which is about 10^{-20} g. For macroscopic spinning bodies, the angular momentum effect on motion has been also been discussed, notably by Mashhoon and Everitt, and the result is generally larger but still beyond the reach of presently contemplated EP tests [34].

IX. SUMMARY AND CONCLUSIONS

We have developed the theory of a spin 1/2 Dirac particle in a Riemann space and its weak field limit in considerable detail. In the low velocity limit for the particle the energies due to the Newtonian or gravitoelectric field and the frame-dragging or gravitomagnetic field take simple and intuitive forms. The small gravitomagnetic effects we have discussed would be quite difficult to detect, but would be of fundamental importance.

The results of the LAGEOS and GP-B experiments and the theoretical results of this paper and Ref. [17] are probably most important in establishing the validity and consistency of general relativity and the gravitomagnetic effects that it implies. Such effects are quite small in earth-based labs and satellite systems, as is clear from (8.3), but as noted in the introduction may play a large role in astrophysical phenomena such as the jets observed in active galactic nuclei, for which the gravitomagnetic fields are much stronger [9].

ACKNOWLEDGMENTS

This work was partially supported by NASA grant 8-39225 to Gravity Probe B and by NSC of Taiwan under Project No. NSC 97-2112-M-002-026-MY3. Pisin Chen thanks Taiwan's National Center for Theoretical Sciences for their support. Thanks go to Robert Wagoner, Francis Everitt, and Alex Silbergleit and other members of the GB-B HEPL Theory Group for useful discussions, and to Mark Kasevich of the Stanford physics department for interesting comments on atomic beam interferometry and equivalence principle experiments. Kung-Yi Su provided valuable help with the manuscript. Elisa Varani thanks Cavallo Pacific for encouragement and support.

APPENDIX A: THE INVERSE DIFFERENTIAL OPERATOR

We briefly study the type of differential operator that appears in (3.7) by solving the differential equation

$$Af + \partial f = (A + \partial)f = F, \quad f = f(x), \quad F = F(x), \quad (\text{A1})$$

where $F(x)$ is a given function that may be expanded as a power series in the region of interest and A is a constant. The solution of the homogeneous equation is

$$f_h = Ce^{-Ax} \quad (C = \text{arbitrary constant}). \quad (\text{A2})$$

The general solution of (A1) is f_h plus any particular solution f_p ; for the particular solution we solve (A1) symbolically as

$$f_p = (A + \partial)^{-1}F = \frac{1}{A} \left(1 - \frac{\partial}{A} + \frac{\partial^2}{A^2} \cdots \right). \quad (\text{A3})$$

It is easily verified that operating with $(A + \partial)$ on the last parenthesis in (A3) does indeed give F .

To further justify the above formal operations we may solve (A1) in a different way. An integrating factor is easily seen to be e^{Ax} , so

$$\partial(e^{Ax}f) = e^{Ax}(A + \partial)f = e^{Ax}F. \quad (\text{A4})$$

Integration then gives the general solution

$$f = e^{-Ax} \int^x e^{-Ax'} F(x') dx' + Ce^{-Ax}. \quad (\text{A5})$$

Since (A1) is linear and F is assumed to be expandable in a power series, we need only consider powers $F = x^n$. Then, we easily evaluate (A5) using integration by parts to obtain

$$f = \frac{1}{A} \left(\frac{x^n}{A} - \frac{nx^{n-1}}{A^2} + \frac{n(n-1)x^{n-2}}{A^3} \cdots + 1 \right) + Ce^{-Ax}. \quad (\text{A6})$$

This agrees with the solutions given in (A2) and (A3).

APPENDIX B: ENERGY-MOMENTUM TENSOR FOR THE DIRAC FIELD

We wish to obtain the energy-momentum tensor for a Dirac field in flatspace, which occurs in (6.8) and (6.9) [18]. We begin with the Lagrangian (3.1) for the free Dirac field and work out the canonical energy-momentum tensor according to the Noether theorem; it is, up to a constant multiplier C ,

$$\begin{aligned} T^{\mu}_{\nu} &= C \left[\frac{\partial L}{\partial \psi_{|\mu}} \psi_{|\nu} + \frac{\partial L}{\partial \bar{\psi}_{|\mu}} \bar{\psi}_{|\nu} - \delta^{\mu}_{\nu} L \right] \\ &= C [a \bar{\psi} i \gamma^{\mu} \psi_{|\nu} - b \bar{\psi}_{|\nu} i \gamma^{\mu} \psi]. \end{aligned} \quad (\text{B1})$$

where we have omitted the term proportional to L since it is zero for a solution of the free Dirac equation. Using the fact that the Dirac and the Klein-Gordon equations are obeyed by ψ , we calculate the two divergences of this tensor to be

$$\begin{aligned} T^{\mu\nu}_{|\mu} &= 0, \\ T^{\mu\nu}_{|\nu} &= C(b - a)[m^2(\bar{\psi} i \gamma^{\mu} \psi) - (\bar{\psi}^{|\nu} i \gamma^{\mu} \psi_{|\nu})]. \end{aligned} \quad (\text{B2})$$

If we choose $b = a$, as in the text, both divergences are zero and the tensor has symmetry in ψ and $\bar{\psi}$. Moreover, we may then consistently symmetrize $T^{\mu\nu}$ and have

$$T^{\mu\nu} = \frac{1}{4} [\bar{\psi} i \gamma^{\mu} \psi^{|\nu} - \bar{\psi}^{|\nu} i \gamma^{\mu} \psi + \bar{\psi} i \gamma^{\nu} \psi^{|\mu} - \bar{\psi}^{|\mu} i \gamma^{\nu} \psi]. \quad (\text{B3})$$

This has now been normalized so that in the low velocity limit

$$T^{00} \approx m \bar{\psi} \psi. \quad (\text{B4})$$

Finally, to include the electromagnetic field we use the minimal substitution recipe $i\partial_\mu \rightarrow i\partial_\mu - eA_\mu$ to get

$$T^{\mu\nu} = \frac{1}{4} [\bar{\psi} i \gamma^\mu \psi^{|\nu} - \bar{\psi}^{|\nu} i \gamma^\mu \psi + \bar{\psi} i \gamma^\nu \psi^{|\mu} - \bar{\psi}^{|\mu} i \gamma^\nu \psi] - \frac{1}{2} [e \bar{\psi} A^\nu \gamma^\mu \psi + e \bar{\psi} A^\mu \gamma^\nu \psi]. \quad (\text{B5})$$

To verify the result (B5) we may calculate the divergence of $T^{\mu\nu}$ to find, after some algebra, that it gives the correct Lorentz force,

$$T^{\mu\nu}{}_{|\mu} = -j_\alpha F^{\mu\alpha} = -(\bar{\psi} \gamma_\alpha \psi) F^{\mu\alpha}. \quad (\text{B6})$$

In the interaction Lagrangian (6.8) the energy-momentum tensor is contracted with the symmetric $h_{\mu\nu}$ so the symmetrization in (B5) is not relevant.

-
- [1] I. Ciufolini *et al.*, *Classical Quantum Gravity* **14**, 2701 (1997); **279**, 2100 (1998); I. Ciufolini and J. A. Wheeler, *Gravitation and Inertia* (Princeton University Press, New Jersey, 1995), Chap. 6; I. Ciufolini *et al.*, in *General Relativity and John Archibald Wheeler*, edited by I. Ciufolini and R. A. Matzner (Springer, Dordrecht, 2010), p. 371.
- [2] R. J. Adler, *The New Physics for the Twenty-first Century*, edited by Gordon Fraser (Cambridge University Press, Cambridge U.K., 2006).
- [3] C. Will, *Theory and Experiment in Gravitational Physics* (Cambridge University Press, Cambridge, U.K., 1981), Chap. 2 and Chap. 5, revised edition 1993.
- [4] H. C. Ohanian and R. Ruffini, *Gravitation and Spacetime* (W. W. Norton, New York, 1976), Chap. 1.
- [5] C. W. F. Everitt *et al.*, *Phys. Rev. Lett.* **106**, 221101 (2011). see also the GP-B <http://einstein.stanford.edu>.
- [6] I. Ciufolini and E. C. Pavlis, *Nature (London), Phys. Sci.* **431**, 958 (2004).
- [7] L. Iorio, H. I. M. Lichtenegger, and C. Corda, *Astrophys. Space Sci.*, **331**, 351 (2010), discuss the phenomenology and accuracy of solar system tests of general relativity.; A criticism of the error analysis of the LAGEOS results is given in L. Iorio, *Europhys. Lett.* **96**, 30001 (2011); A response is given in J. Ries *et al.*, *Europhys. Lett.* **96**, 30002 (2011).
- [8] C. Will, Finally, Results from Gravity Probe B, <http://physics.aps.org/articles/v4/43>.
- [9] K. S. Thorne, *Near Zero: New Frontiers of Physics*, edited by J. D. Fairbank, B. S. Deaver, C. W. F. Everitt, and P. F. Michelson (W. H. Freeman, New York, 1988) p. 573; L. Stella and A. Possenti, *Space Sci. Rev.* **148**, (2010); R. D. Blandford and R. L. Znajek, *Mon. Not. R. Astron. Soc.* **179**, 433 (1977); R. K. Williams, *Phys. Rev.* **51**, No. 10, 5387 (1995).
- [10] S. W. Hawking, *Commun. Math. Phys.* **18**, 301 (1970).
- [11] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge U.K., 1982).
- [12] R. J. Adler, P. Chen, and D. Santiago, *Gen. Relativ. Gravit.* **33**, 2101 (2001).
- [13] V. V. Nesvizhevsky *et al.*, *Nature (London), Phys. Sci.* **415**, 297 (2002).
- [14] S. Dimopoulos, P. W. Graham, J. M. Hogan, and M. A. Kasevich, *Phys. Rev. Lett.* **98**, 111102 (2007); S. Dimopoulos, P. W. Graham, J. M. Hogan, M. A. Kasevich, and S. Rajendran, *Phys. Rev. D* **78**, 12202 (2008).
- [15] S. Weinberg, *Gravitation and Cosmology* (John Wiley, New York, 1972), see Chap. 5 on effects of general relativity on the motion of a classical particle.
- [16] W. Rindler, *Phys. Lett. A* **233**, 25 (1997), and references therein.
- [17] R. J. Adler and P. Chen, *Phys. Rev. D* **82**, 025004 (2010).
- [18] J. D. Bjorken and S. D. Drell, *Relativistic Quantum Mechanics* (McGraw Hill, New York, 1964), we use the conventions and notation for the Dirac equation in flatspace given in Chap. 3; J. D. Bjorken and S. D. Drell, *Relativistic Quantum Fields* (McGraw Hill, New York, 1965), Chap. 13.
- [19] I. Lawrie, *A Unified Grand Tour of Theoretical Physics* (Adam Hilger, Bristol, 1990), generally covariant Dirac theory using tetrads is discussed in Sec. 7.5; see also Weinberg [15] Sec. 12.5.
- [20] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (W. H. Freeman, San Francisco, 1970), Chap. 18; R. J. Adler, M. Bazin, and M. M. Schiffer, *Introduction to General Relativity, 2nd edition* (McGraw Hill, New York, 1975), see Chap. 9.
- [21] R. J. Adler, *Gen. Relativ. Gravit.* **31**, 1837 (1999).
- [22] R. J. Adler and A. S. Silbergleit, *Int. J. Theor. Phys.* **39**, 1291 (2000), note that the definition of the gravitomagnetic field differs by a factor of 2 from our use in this work.
- [23] R. P. Feynman, *Lectures on Gravitation*, lecture notes by F. B. Morinigo and W. C. Wagner (Caltech, Pasadena, 1971); see also Ref. 15, Sec. 6.9; P. C. Peters, *Flat-space Gravitation and Feynman Quantization*, (1965).
- [24] An overview of this vast field is given in D. Oriti, *Approaches to Quantum Gravity* (Cambridge University Press, Cambridge U.K., 2009).
- [25] A. Trautman, in *Encyclopedia of Mathematical Physics*, edited by J. P. Francoise, G. L. Naber, and S. T. Tsou (Elsevier, Oxford U.K., 2006), p. 189; W. T. Ni, *Rep. Prog. Phys.* **73**, 056901 (2010), Secs. 2.2–2.3.
- [26] H. Kleinert, *Electron. J. Theor. Phys.* **24**, 287 (2010).
- [27] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory* (Addison-Wesley, Reading, MA, 1995), Chap. 3.

- [28] R. Shankar, *Principles of Quantum Mechanics* (Plenum, New York, 1994), 2nd ed., Sec. 20.2.
- [29] J. Schwinger, *Am. J. Phys.* **42**, 507 (1974); J. Schwinger, *Gen. Relativ. Gravit.* **7**, 251 (1976).
- [30] R.J. Adler and S.D. Drell, *Phys. Rev. Lett.* **13**, 349 (1964); R. J. Adler, *Phys. Rev.* **141**, 1499 (1966).
- [31] F. W. Hehl and W. T. Ni, *Phys. Rev. D* **42**, 2045 (1990), see Eq. 20.
- [32] R. J. Adler, H. Mueller, and M. L. Perl *Int. J. Mod. Phys. A* **26**, 4959 (2011).
- [33] M. Mathisson, *Acta Phys. Pol.* **6**, No. 218163 (1937); A. Papapetrou, *Proc. R. Soc. A* **209**, 248 (1951); W.G. Dixon, *Phil. Trans. R. Soc. A* **277**, 59 (1974).
- [34] B. Mashhoon, *Classical Quantum Gravity* **17**, 2399 (2000); C.W.F. Everitt, *Proceedings of the 1983 International School on Precision Measurement and Gravity Experiment*, (National Tsing Hua University, Taipei Republic of China, (1983).